

Applications of an operator $H(\alpha, \beta)$ to the Lauricella multivariable hypergeometric functions

A. Hasanov

Institute of Mathematics and Information Technologies, Uzbek Academy of Sciences,

29, F. Hodjaev street, Tashkent 100125, Uzbekistan

E-mail: anvarhasanov@yahoo.com

Abstract

By making use of some techniques based upon certain inverse new pairs of symbolic operators, the author investigate several decomposition formulas associated with Lauricella's hypergeometric functions $F_A^{(r)}, F_B^{(r)}, F_C^{(r)}$ and $F_D^{(r)}$ in r variables. In the three-variable case some of these operational representations are constructed and applied in order to derive the corresponding decomposition formulas when $r = 3$. With the help of these new inverse pairs of symbolic operators, a total 20 decomposition formulas and integral representations are found.

MSC: primary 33C65.

Key Words and Phrases. Decomposition formulas; Lauricella hypergeometric functions; Multiple hypergeometric functions; Generalized hypergeometric functions; Inverse pairs of symbolic operators; Integral representations.

1 Introduction and definitions

A great interest in the theory of hypergeometric functions (that is, hypergeometric functions of several variables) is motivated essentially by the fact that the solutions of many applied problems involving (for example) partial differential equations are obtainable with the help of such hypergeometric function (see, for details, [21, p. 47]; see also the recent works [7-9, 16, 17] and the references cited therein). For instance, the energy absorbed by some nonferromagnetic conductor sphere included in an internal magnetic field can be calculated with the help of such functions [12, 15]. Hypergeometric functions of several variables are used in physical and quantum chemical applications as well [13, 19 and 20]. Especially, many problems in gas dynamics lead to solutions of degenerate second-order partial differential equations, which are then solvable in terms of multiple hypergeometric functions. Among examples, we can cite the problem of adiabatic flat-parallel gas flow without whirlwind, the flow problem of supersonic current from vessel with flat walls, and a number of other problems connected with gas flow [2, 6].

We note that Riemann's functions and fundamental solutions of the degenerate second-order partial differential equations are expressible by means of hypergeometric functions of several variables [7-9]. In investigation of the boundary value problems for these partial differential equations, we need decompositions for hypergeometric functions of several variables in terms of simpler hypergeometric functions of the Gauss and Lauricella types.

Multiple hypergeometric functions (that is, hypergeometric functions in several variables) occur naturally in a wide variety of problems. In particular, the Lauricella functions $F_A^{(r)}$, $F_B^{(r)}$, $F_C^{(r)}$ and $F_D^{(r)}$ in r ($r \in \setminus \{1\}; := \{1, 2, \dots\}$) variables, defined by ([1, p. 114])

$$F_A^{(r)}(\alpha; \beta_1, \dots, \beta_r; \gamma_1, \dots, \gamma_r; x_1, \dots, x_r) = \sum_{m_1, \dots, m_r=0}^{\infty} \frac{(\alpha)_{m_1+\dots+m_r} (\beta_1)_{m_1} \cdots (\beta_r)_{m_r}}{(\gamma_1)_{m_1} \cdots (\gamma_r)_{m_r} m_1! \cdots m_r!} x_1^{m_1} \cdots x_r^{m_r}, \quad (1.1)$$

$$(|x_1| + \cdots + |x_r| < 1),$$

$$F_B^{(r)}(\alpha_1, \dots, \alpha_r; \beta_1, \dots, \beta_r; \gamma; x_1, \dots, x_r) = \sum_{m_1, \dots, m_r=0}^{\infty} \frac{(\alpha_1)_{m_1} \cdots (\alpha_r)_{m_r} (\beta_1)_{m_1} \cdots (\beta_r)_{m_r}}{(\gamma)_{m_1+\dots+m_r} m_1! \cdots m_r!} x_1^{m_1} \cdots x_r^{m_r}, \quad (1.2)$$

$$(\max[|x_1|, \dots, |x_r|] < 1),$$

$$F_C^{(r)}(\alpha, \beta; \gamma_1, \dots, \gamma_r; x_1, \dots, x_r) = \sum_{m_1, \dots, m_r=0}^{\infty} \frac{(\alpha)_{m_1+\dots+m_r} (\beta)_{m_1+\dots+m_r}}{(\gamma_1)_{m_1} \cdots (\gamma_r)_{m_r} m_1! \cdots m_r!} x_1^{m_1} \cdots x_r^{m_r}, \quad (1.3)$$

$$(|\sqrt{x_1}| + \cdots + |\sqrt{x_r}| < 1),$$

$$F_D^{(r)}(\alpha; \beta_1, \dots, \beta_r; \gamma; x_1, \dots, x_r) = \sum_{m_1, \dots, m_r=0}^{\infty} \frac{(\alpha)_{m_1+\dots+m_r} (\beta_1)_{m_1} \cdots (\beta_r)_{m_r}}{(\gamma)_{m_1+\dots+m_r} m_1! \cdots m_r!} x_1^{m_1} \cdots x_r^{m_r}, \quad (1.4)$$

$$(\max[|x_1|, \dots, |x_r|] < 1).$$

Here, and in what follows $(\alpha)_m = \Gamma(\alpha + m) / \Gamma(\alpha)$ denotes the Pochhammer symbol (or the shifted factorial) for all admissible (real or complex) values of α and m .

For various multivariable hypergeometric functions including the Lauricella multivariable functions $F_A^{(r)}$, $F_B^{(r)}$, $F_C^{(r)}$ and $F_D^{(r)}$, defined by (1.1)-(1.4), Hasanov and Srivastava [10, 11] presented a number of decomposition formulas in terms of such simpler hypergeometric functions as the Gauss and Appell functions. The main object of this sequel to the works of Hasanov and Srivastava [10, 11] is to show how some rather elementary techniques based upon certain inverse pairs of symbolic operators would lead us easily to several decomposition formulas associated with Lauricella's hypergeometric function $F_A^{(r)}$, $F_B^{(r)}$, $F_C^{(r)}$ and $F_D^{(r)}$ in r variables and with other multiple hypergeometric functions.

Over six decades ago, Burchnall and Chaundy [3, 4] and Chaundy [5] systematically presented a number of expansion and decomposition formulas for some double hypergeometric functions in series of simpler hypergeometric functions. Their method is based upon the following inverse pairs of symbolic operators:

$$\nabla_{xy}(h) := \frac{\Gamma(h) \Gamma(\delta_1 + \delta_2 + h)}{\Gamma(\delta_1 + h) \Gamma(\delta_2 + h)} = \sum_{k=0}^{\infty} \frac{(-\delta_1)_k (-\delta_2)_k}{(h)_k k!}, \quad (1.5)$$

$$\begin{aligned} \Delta_{xy}(h) &:= \frac{\Gamma(\delta_1 + h) \Gamma(\delta_2 + h)}{\Gamma(h) \Gamma(\delta_1 + \delta_2 + h)} = \sum_{k=0}^{\infty} \frac{(-\delta_1)_k (-\delta_2)_k}{(1 - h - \delta_1 - \delta_2)_k k!} \\ &= \sum_{k=0}^{\infty} \frac{(-1)^k (h)_{2k} (-\delta_1)_k (-\delta_2)_k}{(h + k - 1)_k (h + \delta_1)_k (h + \delta_2)_k k!}, \end{aligned} \quad (1.6)$$

and

$$\begin{aligned} \nabla_{xy}(h) \Delta_{xy}(g) &:= \frac{\Gamma(h) \Gamma(\delta_1 + \delta_2 + h)}{\Gamma(\delta_1 + h) \Gamma(\delta_2 + h)} \frac{\Gamma(\delta_1 + g) \Gamma(\delta_2 + g)}{\Gamma(g) \Gamma(\delta_1 + \delta_2 + g)} \\ &= \sum_{k=0}^{\infty} \frac{(g - h)_k (g)_{2k} (-\delta_1)_k (-\delta_2)_k}{(g + k - 1)_k (g + \delta_1)_k (g + \delta_2)_k k!} \\ &= \sum_{k=0}^{\infty} \frac{(h - g)_k (-\delta_1)_k (-\delta_2)_k}{(h)_k (1 - g - \delta_1 - \delta_2)_k k!} \left(\delta_1 := x \frac{\partial}{\partial x}; \delta_2 := y \frac{\partial}{\partial y} \right). \end{aligned} \quad (1.7)$$

We introduce the following multivariable symbolic operators:

$$H_{x_1, \dots, x_l}(\alpha, \beta) := \frac{\Gamma(\beta) \Gamma(\alpha + \delta_1 + \dots + \delta_l)}{\Gamma(\alpha) \Gamma(\beta + \delta_1 + \dots + \delta_l)} = \sum_{k_1, \dots, k_l=0}^{\infty} \frac{(\beta - \alpha)_{k_1 + \dots + k_l} (-\delta_1)_{k_1} \dots (-\delta_l)_{k_l}}{(\beta)_{k_1 + \dots + k_l} k_1! \dots k_l!} \quad (1.8)$$

and

$$\bar{H}_{x_1, \dots, x_l}(\alpha, \beta) := \frac{\Gamma(\alpha) \Gamma(\beta + \delta_1 + \dots + \delta_l)}{\Gamma(\beta) \Gamma(\alpha + \delta_1 + \dots + \delta_l)} = \sum_{k_1, \dots, k_l=0}^{\infty} \frac{(\beta - \alpha)_{k_1 + \dots + k_l} (-\delta_1)_{k_1} \dots (-\delta_l)_{k_l}}{(1 - \alpha - \delta_1 - \dots - \delta_l)_{k_1 + \dots + k_l} k_1! \dots k_l!}, \quad (1.9)$$

$$\left(\delta_j := x_j \frac{\partial}{\partial x_j}, j = 1, \dots, l; l \in N := \{1, 2, 3, \dots\} \right)$$

where we have applied known multiple hypergeometric summation formulas as follows [1, p. 117]:

$$F_D^{(r)}(\alpha; \beta_1, \dots, \beta_r; \gamma; 1, \dots, 1) = \frac{\Gamma(\gamma) \Gamma(\gamma - \alpha - \beta_1 - \dots - \beta_r)}{\Gamma(\gamma - \alpha) \Gamma(\gamma - \beta_1 - \dots - \beta_r)}$$

$$(\operatorname{Re}(\gamma - \alpha - \beta_1 - \dots - \beta_r) > 0, \gamma \notin N_0^- := \{0, -1, -2, -3, \dots\})$$

for the Lauricella function $F_D^{(r)}$ in r variables, defined by (1.4).

2 Families of decompositions formulas for Lauricella functions

First of all, it is not difficult to derive the following applications of the (multivariable) symbolic operators defined by (1.8) and (1.9):

$$F_A^{(r)}(\alpha; \beta_1, \dots, \beta_r; \gamma_1, \dots, \gamma_r; x_1, \dots, x_r) = H_{x_1, \dots, x_r}(\alpha, \varepsilon) F_A^{(r)}(\varepsilon; \beta_1, \dots, \beta_r, \gamma_1, \dots, \gamma_r; x_1, \dots, x_r), \quad (2.1)$$

$$F_A^{(r)}(\alpha; \beta_1, \dots, \beta_r; \gamma_1, \dots, \gamma_r; x_1, \dots, x_r) = \bar{H}_{x_1, \dots, x_r}(\varepsilon, \alpha) F_A^{(r)}(\varepsilon; \beta_1, \dots, \beta_r, \gamma_1, \dots, \gamma_r; x_1, \dots, x_r), \quad (2.2)$$

$$F_B^{(r)}(\alpha_1, \dots, \alpha_r; \beta_1, \dots, \beta_r, \gamma; x_1, \dots, x_r) = \bar{H}_{x_1, \dots, x_r}(\gamma, \varepsilon) F_B^{(r)}(\alpha_1, \dots, \alpha_r; \beta_1, \dots, \beta_r, \varepsilon; x_1, \dots, x_r), \quad (2.3)$$

$$F_C^{(r)}(\alpha, \beta; \gamma_1, \dots, \gamma_r; x_1, \dots, x_r) = H_{x_1, \dots, x_r}(\alpha, \varepsilon) F_C^{(r)}(\varepsilon, \beta; \gamma_1, \dots, \gamma_r; x_1, \dots, x_r), \quad (2.4)$$

$$F_C^{(r)}(\alpha, \beta; \gamma_1, \dots, \gamma_r; x_1, \dots, x_r) = \bar{H}_{x_1, \dots, x_r}(\varepsilon, \alpha) F_C^{(r)}(\varepsilon, \beta; \gamma_1, \dots, \gamma_r; x_1, \dots, x_r), \quad (2.5)$$

$$F_C^{(r)}(\alpha, \beta; \gamma_1, \dots, \gamma_r; x_1, \dots, x_r) = H_{x_1, \dots, x_r}(\alpha, \varepsilon_1) H_{x_1, \dots, x_r}(\beta, \varepsilon_2) F_C^{(r)}(\varepsilon_1, \varepsilon_2; \gamma_1, \dots, \gamma_r; x_1, \dots, x_r), \quad (2.6)$$

$$F_C^{(r)}(\alpha, \beta; \gamma_1, \dots, \gamma_r; x_1, \dots, x_r) = \bar{H}_{x_1, \dots, x_r}(\varepsilon_1, \alpha) \bar{H}_{x_1, \dots, x_r}(\varepsilon_2, \beta) F_C^{(r)}(\varepsilon_1, \varepsilon_2; \gamma_1, \dots, \gamma_r; x_1, \dots, x_r), \quad (2.7)$$

$$F_D^{(r)}(\alpha; \beta_1, \dots, \beta_r; \gamma; x_1, \dots, x_r) = H_{x_1, \dots, x_r}(\alpha, \varepsilon) F_D^{(r)}(\varepsilon; \beta_1, \dots, \beta_r; \gamma; x_1, \dots, x_r), \quad (2.8)$$

$$F_D^{(r)}(\alpha; \beta_1, \dots, \beta_r; \gamma; x_1, \dots, x_r) = \bar{H}_{x_1, \dots, x_r}(\varepsilon, \alpha) F_D^{(r)}(\varepsilon; \beta_1, \dots, \beta_r; \gamma; x_1, \dots, x_r), \quad (2.9)$$

$$F_D^{(r)}(\alpha; \beta_1, \dots, \beta_r; \gamma; x_1, \dots, x_r) = H_{x_1, \dots, x_r}(\varepsilon, \gamma) F_D^{(r)}(\alpha; \beta_1, \dots, \beta_r; \varepsilon; x_1, \dots, x_r), \quad (2.10)$$

$$F_D^{(r)}(\alpha; \beta_1, \dots, \beta_r; \gamma; x_1, \dots, x_r) = H_{x_1, \dots, x_r}(\alpha, \gamma) (1 - x_1)^{-\beta_1} \dots (1 - x_r)^{-\beta_r}, \quad (2.11)$$

$$(1 - x_1)^{-\beta_1} \dots (1 - x_r)^{-\beta_r} = \bar{H}_{x_1, \dots, x_r}(\alpha, \gamma) F_D^{(r)}(\alpha; \beta_1, \dots, \beta_r; \gamma; x_1, \dots, x_r). \quad (2.12)$$

In the operator identities (2.6) and (2.7), the superposition of the operators involved can be represented as follows:

$$H_{x_1, \dots, x_r}(\alpha, \varepsilon_1) H_{x_1, \dots, x_r}(\beta, \varepsilon_2)$$

$$= \sum_{k_1, \dots, k_r, l_1, \dots, l_r=0}^{\infty} \frac{(\varepsilon_1 - \alpha)_{k_1 + \dots + k_r} (\varepsilon_2 - \beta)_{l_1 + \dots + l_r} (\beta)_{k_1 + \dots + k_r} (-\delta_1)_{k_1 + l_1} \dots (-\delta_r)_{k_r + l_r}}{(\varepsilon_1)_{k_1 + \dots + k_r} (\varepsilon_2)_{l_1 + \dots + l_r} k_1! \dots k_r! l_1! \dots l_r!} \quad (2.13)$$

and

$$\begin{aligned}
& \bar{H}_{x_1, \dots, x_r}(\varepsilon_1, \alpha) \bar{H}_{x_1, \dots, x_r}(\varepsilon_2, \beta) \\
&= \sum_{k_1, \dots, k_r, l_1, \dots, l_r=0}^{\infty} \frac{(-1)^{k_1+\dots+k_r} (\alpha - \varepsilon_1)_{k_1+\dots+k_r} (\beta - \varepsilon_2)_{k_1+\dots+k_r+l_1+\dots+l_r} (\beta)_{k_1+\dots+k_r}}{(\beta - \varepsilon_2)_{k_1+\dots+k_r} k_1! \dots k_r! l_1! \dots l_r!} \\
&\times \frac{(-\delta_1)_{k_1+l_1} \dots (-\delta_r)_{k_r+l_r}}{(1 - \varepsilon_1 - \delta_1 - \dots - \delta_r)_{k_1+\dots+k_r} (1 - \varepsilon_2 - \delta_1 - \dots - \delta_r)_{k_1+\dots+k_r+l_1+\dots+l_r}}.
\end{aligned} \tag{2.14}$$

In view of the know Mellin-Barnes contour integral representations for the Lauricella functions $F_A^{(r)}$, $F_B^{(r)}$, $F_C^{(r)}$ and $F_D^{(r)}$, it is not difficult to give alternative proofs of the operator identities (2.1) to (2.12) above by using the Mellin and the inverse Mellin transformations (see, for example, [14]). The details involved in these alternative derivations of the operator identities (2.1) to (2.12) are being omitted here.

By virtue of the derivative formulas for the Lauricella functions, and also of some standard properties of hypergeometric functions, we find each of the following decompositions formulas for the Lauricella functions $F_A^{(r)}$, $F_B^{(r)}$, $F_C^{(r)}$ and $F_D^{(r)}$:

$$\begin{aligned}
& F_A^{(r)}(\alpha; \beta_1, \dots, \beta_r; \gamma_1, \dots, \gamma_r; x_1, \dots, x_r) = \sum_{k_1, \dots, k_r=0}^{\infty} \frac{(-1)^{k_1+\dots+k_r} (\varepsilon - \alpha)_{k_1+\dots+k_r} (\beta_1)_{k_1} \dots (\beta_r)_{k_r}}{(\gamma_1)_{k_1} \dots (\gamma_r)_{k_r} k_1! \dots k_r!} \\
& \times x_1^{k_1} x_2^{k_2} \dots x_r^{k_r} F_A^{(r)}(\varepsilon + k_1 + \dots + k_r; \beta_1 + k_1, \dots, \beta_r + k_r; \gamma_1 + k_1, \dots, \gamma_r + k_r; x_1, \dots, x_r),
\end{aligned} \tag{2.15}$$

$$\begin{aligned}
& F_A^{(r)}(\alpha; \beta_1, \dots, \beta_r; \gamma_1, \dots, \gamma_r; x_1, \dots, x_r) = \sum_{k_1, \dots, k_r=0}^{\infty} \frac{(\alpha - \varepsilon)_{k_1+\dots+k_r} (\beta_1)_{k_1} (\beta_2)_{k_2} \dots (\beta_r)_{k_r}}{(\gamma_1)_{k_1} (\gamma_2)_{k_2} \dots (\gamma_r)_{k_r} k_1! \dots k_r!} \\
& \times x_1^{k_1} \dots x_r^{k_r} F_A^{(r)}(\varepsilon; \beta_1 + k_1, \dots, \beta_r + k_r; \gamma_1 + k_1, \dots, \gamma_r + k_r; x_1, \dots, x_r),
\end{aligned} \tag{2.16}$$

$$\begin{aligned}
& F_B^{(r)}(\alpha_1, \dots, \alpha_r; \beta_1, \dots, \beta_r; \gamma; x_1, \dots, x_r) \\
&= \sum_{k_1, \dots, k_r=0}^{\infty} \frac{(-1)^{k_1+\dots+k_r} (\gamma - \varepsilon)_{k_1+\dots+k_r} (\alpha_1)_{k_1} \dots (\alpha_r)_{k_r} (\beta_1)_{k_1} \dots (\beta_r)_{k_r}}{(\gamma)_{k_1+\dots+k_r} (\varepsilon)_{k_1+\dots+k_r} k_1! \dots k_r!} \\
& \times x_1^{k_1} \dots x_r^{k_r} F_B^{(r)}(\alpha_1 + k_1, \dots, \alpha_r + k_r; \beta_1 + k_1, \dots, \beta_r + k_r; \varepsilon + k_1 + \dots + k_r; x_1, \dots, x_r),
\end{aligned} \tag{2.17}$$

$$\begin{aligned}
& F_C^{(r)}(\alpha, \beta; \gamma_1, \dots, \gamma_r; x_1, \dots, x_r) = \sum_{k_1, \dots, k_r=0}^{\infty} \frac{(-1)^{k_1+\dots+k_r} (\varepsilon - \alpha)_{k_1+\dots+k_r} (\beta)_{k_1+\dots+k_r}}{(\gamma_1)_{k_1} \dots (\gamma_r)_{k_r} k_1! \dots k_r!} x_1^{k_1} \dots x_r^{k_r} \\
& \times F_C^{(r)}(\varepsilon + k_1 + \dots + k_r, \beta + k_1 + \dots + k_r; \gamma_1 + k_1, \dots, \gamma_r + k_r; x_1, \dots, x_r),
\end{aligned} \tag{2.18}$$

$$\begin{aligned}
& F_C^{(r)}(\alpha, \beta; \gamma_1, \dots, \gamma_r; x_1, \dots, x_r) = \sum_{k_1, \dots, k_r=0}^{\infty} \frac{(\alpha - \varepsilon)_{k_1+\dots+k_r} (\beta)_{k_1+\dots+k_r}}{(\gamma_1)_{k_1} \dots (\gamma_r)_{k_r} k_1! \dots k_r!} x_1^{k_1} \dots x_r^{k_r} \\
& \times F_C^{(r)}(\varepsilon, \beta + k_1 + \dots + k_r; \gamma_1 + k_1, \dots, \gamma_r + k_r; x_1, \dots, x_r),
\end{aligned} \tag{2.19}$$

$$\begin{aligned}
& F_C^{(r)}(\alpha, \beta; \gamma_1, \dots, \gamma_r; x_1, \dots, x_r) = \sum_{k_1, \dots, k_r, l_1, \dots, l_r=0}^{\infty} \frac{(-1)^{k_1+\dots+k_r+l_1+\dots+l_r} (\varepsilon_1 - \alpha)_{k_1+\dots+k_r}}{(\varepsilon_1)_{k_1+\dots+k_r} (\gamma_1)_{k_1+l_1} \dots (\gamma_r)_{k_r+l_r}} \\
& \frac{(\varepsilon_2 - \beta)_{l_1+\dots+l_r} (\beta)_{k_1+\dots+k_r} (\varepsilon_1)_{k_1+\dots+k_r+l_1+\dots+l_r}}{k_1! \dots k_r! l_1! \dots l_r!} x_1^{k_1+l_1} \dots x_r^{k_r+l_r} \\
& \times F_C^{(r)}(\varepsilon_1 + k_1 + \dots + k_r + l_1 + \dots + l_r, \varepsilon_2 + k_1 + \dots + k_r + l_1 + \dots + l_r; \\
& \gamma_1 + k_1 + l_1, \dots, \gamma_r + k_r + l_r; x_1, \dots, x_r),
\end{aligned} \tag{2.20}$$

$$\begin{aligned}
& F_C^{(r)}(\alpha, \beta; \gamma_1, \dots, \gamma_r; x_1, \dots, x_r) \\
&= \sum_{k_1, \dots, k_r, l_1, \dots, l_r=0}^{\infty} \frac{(-1)^{l_1+\dots+l_r} (\alpha - \varepsilon_1)_{k_1+\dots+k_r} (\beta - \varepsilon_2)_{k_1+\dots+k_r+l_1+\dots+l_r} (\beta)_{k_1+\dots+k_r}}{(\beta - \varepsilon_2)_{k_1+\dots+k_r} (\gamma_1)_{k_1+l_1} \dots (\gamma_r)_{k_r+l_r} k_1! \dots k_r! l_1! \dots l_r!} x_1^{k_1+l_1} \dots x_r^{k_r+l_r} \\
&\times F_C^{(r)}(\varepsilon_1 + l_1 + \dots + l_r, \varepsilon_2 + l_1 + \dots + l_r; \gamma_1 + k_1 + l_1, \dots, \gamma_r + k_r + l_r; x_1, \dots, x_r), \tag{2.21}
\end{aligned}$$

$$\begin{aligned}
& F_D^{(r)}(\alpha; \beta_1, \dots, \beta_r; \gamma; x_1, \dots, x_r) = \sum_{k_1, \dots, k_r=0}^{\infty} \frac{(-1)^{k_1+\dots+k_r} (\varepsilon - \alpha)_{k_1+\dots+k_r} (\beta_1)_{k_1} \dots (\beta_r)_{k_r}}{(\gamma)_{k_1+\dots+k_r} k_1! \dots k_r!} x_1^{k_1} \dots x_r^{k_r} \\
&\times F_D^{(r)}(\varepsilon + k_1 + \dots + k_r; \beta_1 + k_1, \dots, \beta_r + k_r; \gamma + k_1 + \dots + k_r; x_1, \dots, x_r), \tag{2.22}
\end{aligned}$$

$$\begin{aligned}
& F_D^{(r)}(\alpha; \beta_1, \dots, \beta_r; \gamma; x_1, \dots, x_r) = \sum_{k_1, \dots, k_r=0}^{\infty} \frac{(\alpha - \varepsilon)_{k_1+\dots+k_r} (\beta_1)_{k_1} \dots (\beta_r)_{k_r}}{(\gamma)_{k_1+\dots+k_r} k_1! \dots k_r!} x_1^{k_1} \dots x_r^{k_r} \\
&\times F_D^{(r)}(\varepsilon; \beta_1 + k_1, \dots, \beta_r + k_r; \gamma + k_1 + \dots + k_r; x_1, \dots, x_r), \tag{2.23}
\end{aligned}$$

$$\begin{aligned}
& F_D^{(r)}(\alpha; \beta_1, \dots, \beta_r; \gamma; x_1, \dots, x_r) \\
&= \sum_{k_1, \dots, k_r=0}^{\infty} \frac{(-1)^{k_1+\dots+k_r} (\gamma - \varepsilon)_{k_1+\dots+k_r} (\alpha)_{k_1+\dots+k_r} (\beta_1)_{k_1} \dots (\beta_r)_{k_r}}{(\gamma)_{k_1+\dots+k_r} (\varepsilon)_{k_1+\dots+k_r} k_1! \dots k_r!} x_1^{k_1} \dots x_r^{k_r} \\
&\times F_D^{(r)}(\alpha + k_1 + \dots + k_r; \beta_1 + k_1, \dots, \beta_r + k_r; \varepsilon + k_1 + \dots + k_r; x_1, \dots, x_r), \tag{2.24}
\end{aligned}$$

$$\begin{aligned}
& F_D^{(r)}(\alpha; \beta_1, \dots, \beta_r; \gamma; x_1, \dots, x_r) \\
&= (1 - x_1)^{-\beta_1} \dots (1 - x_r)^{-\beta_r} F_D^{(r)}\left(\gamma - \alpha; \beta_1, \dots, \beta_r; \gamma; \frac{x_1}{x_1 - 1}, \dots, \frac{x_r}{x_r - 1}\right), \tag{2.25}
\end{aligned}$$

$$\begin{aligned}
(1 - x_1)^{-\beta_1} \dots (1 - x_r)^{-\beta_r} &= \sum_{k_1, \dots, k_r=0}^{\infty} \frac{(\gamma - \alpha)_{k_1+\dots+k_r} (\beta_1)_{k_1} \dots (\beta_r)_{k_r}}{(\gamma)_{k_1+\dots+k_r} k_1! \dots k_r!} x_1^{k_1} \dots x_r^{k_r} \\
&\times F_D^{(r)}(\alpha; \beta_1 + k_1, \dots, \beta_r + k_r; \gamma + k_1 + \dots + k_r; x_1, \dots, x_r), \tag{2.26}
\end{aligned}$$

Our operational derivations of the decomposition formulas (2.15) to (2.26) would indeed run parallel to those presented in the earlier works, which we have already cited in the preceding sections. In addition to the various operator expressions, we also make use of the following operator identities [18, p. 93]:

$$(\delta + \alpha)_n \{f(\xi)\} = \xi^{1-\alpha} \frac{d^n}{d\xi^n} \{\xi^{\alpha+n-1} f(\xi)\} \tag{2.27}$$

$$\left(\delta := \xi \frac{d}{d\xi}; \alpha \in; n \in_0 := \cup \{0\}; := \{1, 2, 3, \dots\} \right)$$

and

$$(-\delta)_n \{f(\xi)\} = (-1)^n \xi^n \frac{d^n}{d\xi^n} \{f(\xi)\}, \left(\delta := \xi \frac{d}{d\xi}; n \in_0 \right) \tag{2.28}$$

for every analytic function $f(\xi)$. Many other analogous decomposition formulas can similarly be derived for the Lauricella functions $F_A^{(r)}$, $F_B^{(r)}$, $F_C^{(r)}$ and $F_D^{(r)}$ in r variables, but with various different parametric constraints.

3 Basic operator identities for Lauricella functions in the three-variable cases

In this section, we consider the Lauricella functions $F_A^{(3)}$, $F_B^{(3)}$, $F_C^{(3)}$ and $F_D^{(3)}$ in three variables. We reiterate the aforementioned fact that numerous other analogous decompositions formulas can similarly

be derived for the Lauricella triple hypergeometric functions $F_A^{(3)}$, $F_B^{(3)}$, $F_C^{(3)}$ and $F_D^{(3)}$ with different parametric constrains. For example, each of the following operational representations would lead us a decomposition formula for a special $F_A^{(3)}$, $F_B^{(3)}$, $F_C^{(3)}$ and $F_D^{(3)}$ in three variables.

$$F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) = H_x(\beta_1, \varepsilon_1) F_A(\alpha; \varepsilon_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z), \quad (3.1)$$

$$F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) = \bar{H}_x(\varepsilon_1, \beta_1) F_A(\alpha; \varepsilon_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z), \quad (3.2)$$

$$F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) = H_x(\beta_1, \varepsilon_1) H_y(\beta_2, \varepsilon_2) F_A(\alpha; \varepsilon_1, \varepsilon_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z), \quad (3.3)$$

$$F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) = \bar{H}_x(\varepsilon_1, \beta_1) \bar{H}_y(\varepsilon_2, \beta_2) F_A(\alpha; \varepsilon_1, \varepsilon_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z), \quad (3.4)$$

$$F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) = H_x(\beta_1, \varepsilon_1) H_y(\beta_2, \varepsilon_2) H_z(\beta_3, \varepsilon_3) F_A(\alpha; \varepsilon_1, \varepsilon_2, \varepsilon_3; \gamma_1, \gamma_2, \gamma_3; x, y, z), \quad (3.5)$$

$$F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) = \bar{H}_x(\varepsilon_1, \beta_1) \bar{H}_y(\varepsilon_2, \beta_2) \bar{H}_z(\varepsilon_3, \beta_3) F_A(\alpha; \varepsilon_1, \varepsilon_2, \varepsilon_3; \gamma_1, \gamma_2, \gamma_3; x, y, z), \quad (3.6)$$

$$F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) = H_x(\beta_1, \gamma_1) (1-x)^{-\alpha} F_2\left(\alpha; \beta_2, \beta_3; \gamma_2, \gamma_3; \frac{y}{1-x}, \frac{z}{1-x}\right), \quad (3.7)$$

$$(1-x)^{-\alpha} F_2\left(\alpha; \beta_2, \beta_3; \gamma_2, \gamma_3; \frac{y}{1-x}, \frac{z}{1-x}\right) = \bar{H}_x(\beta_1, \gamma_1) F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z), \quad (3.8)$$

$$F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) = H_x(\beta_1, \gamma_1) H_y(\beta_2, \gamma_2) (1-x-y)^{-\alpha} F\left(\alpha, \beta_3; \gamma_3; \frac{z}{1-x-y}\right), \quad (3.9)$$

$$(1-x-y)^{-\alpha} F\left(\alpha, \beta_3; \gamma_3; \frac{z}{1-x-y}\right) = \bar{H}_x(\beta_1, \gamma_1) \bar{H}_y(\beta_2, \gamma_2) F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z), \quad (3.10)$$

$$F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) = H_x(\beta_1, \gamma_1) H_y(\beta_2, \gamma_2) H_z(\beta_3, \gamma_3) (1-x-y-z)^{-\alpha}, \quad (3.11)$$

$$(1-x-y-z)^{-\alpha} = \bar{H}_x(\beta_1, \gamma_1) \bar{H}_y(\beta_2, \gamma_2) \bar{H}_z(\beta_3, \gamma_3) F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z), \quad (3.12)$$

$$F_B(\alpha_1, \alpha_2, \alpha_3; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = H_x(\alpha_1, \varepsilon_1) F_B(\varepsilon_1, \alpha_2, \alpha_3; \beta_1, \beta_2, \beta_3; \gamma; x, y, z), \quad (3.13)$$

$$F_B(\alpha_1, \alpha_2, \alpha_3; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = \bar{H}_x(\varepsilon_1, \alpha_1) F_B(\varepsilon_1, \alpha_2, \alpha_3; \beta_1, \beta_2, \beta_3; \gamma; x, y, z), \quad (3.14)$$

$$F_B(\alpha_1, \alpha_2, \alpha_3; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = H_x(\alpha_1, \varepsilon_1) H_y(\alpha_2, \varepsilon_2) F_B(\varepsilon_1, \varepsilon_2, \alpha_3; \beta_1, \beta_2, \beta_3; \gamma; x, y, z), \quad (3.15)$$

$$F_B(\alpha_1, \alpha_2, \alpha_3; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = \bar{H}_x(\varepsilon_1, \alpha_1) \bar{H}_y(\varepsilon_2, \alpha_2) F_B(\varepsilon_1, \varepsilon_2, \alpha_3; \beta_1, \beta_2, \beta_3; \gamma; x, y, z), \quad (3.16)$$

$$F_B(\alpha_1, \alpha_2, \alpha_3; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = H_x(\alpha_1, \varepsilon_1) H_y(\alpha_2, \varepsilon_2) H_z(\alpha_3, \varepsilon_3) F_B(\varepsilon_1, \varepsilon_2, \varepsilon_3; \beta_1, \beta_2, \beta_3; \gamma; x, y, z), \quad (3.17)$$

$$F_B(\alpha_1, \alpha_2, \alpha_3; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = \bar{H}_x(\varepsilon_1, \alpha_1) \bar{H}_y(\varepsilon_2, \alpha_2) \bar{H}_z(\varepsilon_3, \alpha_3) F_B(\varepsilon_1, \varepsilon_2, \varepsilon_3; \beta_1, \beta_2, \beta_3; \gamma; x, y, z), \quad (3.18)$$

$$F_C(\alpha, \beta; \gamma_1, \gamma_2, \gamma_3; x, y, z) = H_x(\varepsilon_1, \gamma_1) F_C(\alpha, \beta; \varepsilon_1, \gamma_2, \gamma_3; x, y, z), \quad (3.19)$$

$$F_C(\alpha, \beta; \gamma_1, \gamma_2, \gamma_3; x, y, z) = H_x(\varepsilon_1, \gamma_1) H_y(\varepsilon_2, \gamma_2) F_C(\alpha, \beta; \varepsilon_1, \varepsilon_2, \gamma_3; x, y, z), \quad (3.20)$$

$$F_C(\alpha, \beta; \gamma_1, \gamma_2, \gamma_3; x, y, z) = H_x(\varepsilon_1, \gamma_1) H_y(\varepsilon_2, \gamma_2) H_z(\varepsilon_3, \gamma_3) F_C(\alpha, \beta; \varepsilon_1, \varepsilon_2, \varepsilon_3; x, y, z), \quad (3.21)$$

$$F_D(\alpha; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = H_x(\beta_1, \varepsilon_1) F_D(\alpha; \varepsilon_1, \beta_2, \beta_3; \gamma; x, y, z), \quad (3.22)$$

$$F_D(\alpha; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = \bar{H}_x(\varepsilon_1, \beta_1) F_D(\alpha; \varepsilon_1, \beta_2, \beta_3; \gamma; x, y, z), \quad (3.23)$$

$$F_D(\alpha; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = H_x(\beta_1, \varepsilon_1) H_y(\beta_2, \varepsilon_2) F_D(\alpha; \varepsilon_1, \varepsilon_2, \beta_3; \gamma; x, y, z), \quad (3.24)$$

$$F_D(\alpha; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = \bar{H}_x(\varepsilon_1, \beta_1) \bar{H}_y(\varepsilon_2, \beta_2) F_D(\alpha; \varepsilon_1, \varepsilon_2, \beta_3; \gamma; x, y, z), \quad (3.25)$$

$$F_D(\alpha; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = H_x(\beta_1, \varepsilon_1) H_y(\beta_2, \varepsilon_2) H_z(\beta_3, \varepsilon_3) F_D(\alpha; \varepsilon_1, \varepsilon_2, \varepsilon_3; \gamma; x, y, z), \quad (3.26)$$

$$F_D(\alpha; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = \bar{H}_x(\varepsilon_1, \beta_1) \bar{H}_y(\varepsilon_2, \beta_2) \bar{H}_z(\varepsilon_3, \beta_3) F_D(\alpha; \varepsilon_1, \varepsilon_2, \varepsilon_3; \gamma; x, y, z), \quad (3.27)$$

where, for notational convenience

$$F_A := F_A^{(3)}, F_B := F_B^{(3)}, F_C := F_C^{(3)}, F_C := F_C^{(3)}, F_2 := F_A^{(2)}. \quad (3.28)$$

By virtue of the derivative formulas for the Lauricella functions, and also of some standard properties of hypergeometric functions, we find each of the following decompositions formulas for the Lauricella functions $F_A^{(3)}$, $F_B^{(3)}$, $F_C^{(3)}$ and $F_D^{(3)}$:

$$\begin{aligned} F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) \\ = \sum_{i=0}^{\infty} \frac{(-1)^i (\varepsilon_1 - \beta_1)_i (\alpha)_i}{(\gamma_1)_i i!} x^i F_A(\alpha + i; \varepsilon_1 + i, \beta_2, \beta_3; \gamma_1 + i, \gamma_2, \gamma_3; x, y, z), \end{aligned} \quad (3.29)$$

$$F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) = \sum_{i=0}^{\infty} \frac{(\beta_1 - \varepsilon_1)_i (\alpha)_i}{(\gamma_1)_i i!} x^i F_A(\alpha + i; \varepsilon_1, \beta_2, \beta_3; \gamma_1 + i, \gamma_2, \gamma_3; x, y, z), \quad (3.30)$$

$$\begin{aligned} F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) = \sum_{i,j=0}^{\infty} \frac{(-1)^{i+j} (\alpha)_{i+j} (\varepsilon_1 - \beta_1)_i (\varepsilon_2 - \beta_2)_j}{(\gamma_1)_i (\gamma_2)_j i! j!} x^i y^j \\ \times F_A(\alpha + i + j; \varepsilon_1 + i, \varepsilon_2 + j, \beta_3; \gamma_1 + i, \gamma_2 + j, \gamma_3; x, y, z), \end{aligned} \quad (3.31)$$

$$\begin{aligned} F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) = \sum_{i,j=0}^{\infty} \frac{(\alpha)_{i+j} (\beta_1 - \varepsilon_1)_i (\beta_2 - \varepsilon_2)_j}{(\gamma_1)_i (\gamma_2)_j i! j!} x^i y^j \\ \times F_A(\alpha + i + j; \varepsilon_1, \varepsilon_2, \beta_3; \gamma_1 + i, \gamma_2 + j, \gamma_3; x, y, z), \end{aligned} \quad (3.32)$$

$$\begin{aligned} F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) \\ = \sum_{i,j,k=0}^{\infty} \frac{(-1)^{i+j+k} (\alpha)_{i+j+k} (\varepsilon_1 - \beta_1)_i (\varepsilon_2 - \beta_2)_j (\varepsilon_3 - \beta_3)_k}{(\gamma_1)_i (\gamma_2)_j (\gamma_3)_k i! j! k!} x^i y^j z^k \\ \times F_A(\alpha + i + j + k; \varepsilon_1 + i, \varepsilon_2 + j, \varepsilon_3 + k; \gamma_1 + i, \gamma_2 + j, \gamma_3 + k; x, y, z), \end{aligned} \quad (3.33)$$

$$\begin{aligned} F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) = \sum_{i,j,k=0}^{\infty} \frac{(\alpha)_{i+j+k} (\beta_1 - \varepsilon_1)_i (\beta_2 - \varepsilon_2)_j (\beta_3 - \varepsilon_3)_k}{(\gamma_1)_i (\gamma_2)_j (\gamma_3)_k i! j! k!} x^i y^j z^k \\ \times F_A(\alpha + i + j + k; \varepsilon_1, \varepsilon_2, \varepsilon_3; \gamma_1 + i, \gamma_2 + j, \gamma_3 + k; x, y, z), \end{aligned} \quad (3.34)$$

$$\begin{aligned} F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) \\ = (1-x)^{-\alpha} \sum_{i=0}^{\infty} \frac{(\alpha)_i (\gamma_1 - \beta_1)_i}{(\gamma_1)_i i!} \left(\frac{x}{x-1} \right)^i F_2 \left(\alpha + i; \beta_2, \beta_3; \gamma_2, \gamma_3; \frac{y}{1-x}, \frac{z}{1-x} \right), \end{aligned} \quad (3.35)$$

$$\begin{aligned} (1-x)^{-\alpha} F_2 \left(\alpha; \beta_2, \beta_3; \gamma_2, \gamma_3; \frac{y}{1-x}, \frac{z}{1-x} \right) \\ = \sum_{i=0}^{\infty} \frac{(\alpha)_i (\gamma_1 - \beta_1)_i}{(\gamma_1)_i i!} x^i F_A(\alpha + i; \beta_1, \beta_2, \beta_3; \gamma_1 + i, \gamma_2, \gamma_3; x, y, z), \end{aligned} \quad (3.36)$$

$$F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) = (1-x-y)^{-\alpha} \sum_{i,j=0}^{\infty} \frac{(-1)^{i+j} (\alpha)_{i+j} (\gamma_1 - \beta_1)_i (\gamma_2 - \beta_2)_j}{(\gamma_1)_i (\gamma_2)_j i! j!} \times \left(\frac{x}{1-x-y} \right)^i \left(\frac{y}{1-x-y} \right)^j F\left(\alpha + i + j, \beta_3; \gamma_3; \frac{z}{1-x-y}\right), \quad (3.37)$$

$$(1-x-y)^{-\alpha} F\left(\alpha, \beta_3; \gamma_3; \frac{z}{1-x-y}\right) = \sum_{i,j=0}^{\infty} \frac{(\alpha)_{i+j} (\gamma_1 - \beta_1)_i (\gamma_2 - \beta_2)_j}{(\gamma_1)_i (\gamma_2)_j i! j!} x^i y^j F_A(\alpha + i + j; \beta_1, \beta_2, \beta_3; \gamma_1 + i, \gamma_2 + j, \gamma_3; x, y, z), \quad (3.38)$$

$$F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) = (1-x-y-z)^{-\alpha} \times F_A\left(\alpha; \gamma_1 - \beta_1, \gamma_2 - \beta_2, \gamma_3 - \beta_3; \gamma_1, \gamma_2, \gamma_3; \frac{x}{x+y+z-1}, \frac{y}{x+y+z-1}, \frac{z}{x+y+z-1}\right), \quad (3.39)$$

$$(1-x-y-z)^{-\alpha} = \sum_{i,j,k=0}^{\infty} \frac{(\alpha)_{i+j+k} (\gamma_1 - \beta_1)_i (\gamma_2 - \beta_2)_j (\gamma_3 - \beta_3)_k}{(\gamma_1)_i (\gamma_2)_j (\gamma_3)_k i! j! k!} x^i y^j z^k \times F_A(\alpha + i + j + k; \beta_1, \beta_2, \beta_3; \gamma_1 + i, \gamma_2 + j, \gamma_3 + k; x, y, z), \quad (3.40)$$

$$F_B(\alpha_1, \alpha_2, \alpha_3; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = \sum_{i=0}^{\infty} \frac{(-1)^i (\varepsilon_1 - \alpha_1)_i (\beta_1)_i}{(\gamma)_i i!} x^i F_B(\varepsilon_1 + i, \alpha_2, \alpha_3; \beta_1 + i, \beta_2, \beta_3; \gamma + i; x, y, z), \quad (3.41)$$

$$F_B(\alpha_1, \alpha_2, \alpha_3; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = \sum_{i=0}^{\infty} \frac{(\alpha_1 - \varepsilon_1)_i (\beta_1)_i}{(\gamma_1)_i i!} x^i F_B(\varepsilon_1, \alpha_2, \alpha_3; \beta_1 + i, \beta_2, \beta_3; \gamma + i; x, y, z), \quad (3.42)$$

$$F_B(\alpha_1, \alpha_2, \alpha_3; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = \sum_{i,j=0}^{\infty} \frac{(-1)^{i+j} (\varepsilon_1 - \alpha_1)_i (\varepsilon_2 - \alpha_2)_j (\beta_1)_i (\beta_2)_j}{(\gamma)_{i+j} i! j!} x^i y^j \times F_B(\varepsilon_1 + i, \varepsilon_2 + j, \alpha_3; \beta_1 + i, \beta_2 + j, \beta_3; \gamma + i + j; x, y, z), \quad (3.43)$$

$$F_B(\alpha_1, \alpha_2, \alpha_3; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = \sum_{i,j=0}^{\infty} \frac{(\alpha_1 - \varepsilon_1)_i (\alpha_2 - \varepsilon_2)_i (\beta_1)_i (\beta_2)_j}{(\gamma_1)_{i+j} i! j!} x^i y^j F_B(\varepsilon_1, \varepsilon_2, \alpha_3; \beta_1 + i, \beta_2 + j, \beta_3; \gamma + i + j; x, y, z), \quad (3.44)$$

$$F_B(\alpha_1, \alpha_2, \alpha_3; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = \sum_{i,j,k=0}^{\infty} \frac{(-1)^{i+j+k} (\varepsilon_1 - \alpha_1)_i (\varepsilon_2 - \alpha_2)_j (\varepsilon_3 - \alpha_3)_k (\beta_1)_i (\beta_2)_j (\beta_3)_k}{(\gamma)_{i+j+k} i! j! k!} x^i y^j z^k \times F_B(\varepsilon_1 + i, \varepsilon_2 + j, \varepsilon_3 + k; \beta_1 + i, \beta_2 + j, \beta_3 + k; \gamma + i + j + k; x, y, z), \quad (3.45)$$

$$F_B(\alpha_1, \alpha_2, \alpha_3; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = \sum_{i,j,k=0}^{\infty} \frac{(\alpha_1 - \varepsilon_1)_i (\alpha_2 - \varepsilon_2)_i (\alpha_3 - \varepsilon_3)_k (\beta_1)_i (\beta_2)_j (\beta_3)_k}{(\gamma_1)_{i+j+k} i! j! k!} x^i y^j z^k \times F_B(\varepsilon_1, \varepsilon_2, \varepsilon_3; \beta_1 + i, \beta_2 + j, \beta_3 + k; \gamma + i + j + k; x, y, z), \quad (3.46)$$

$$F_C(\alpha, \beta; \gamma_1, \gamma_2, \gamma_3; x, y, z) = \sum_{i=0}^{\infty} \frac{(-1)^i (\alpha)_i (\beta)_i (\gamma_1 - \varepsilon_1)_i}{(\gamma_1)_i (\varepsilon_1)_i i!} x^i F_C(\alpha + i, \beta + i; \varepsilon_1 + i, \gamma_2, \gamma_3; x, y, z), \quad (3.47)$$

$$F_C(\alpha, \beta; \gamma_1, \gamma_2, \gamma_3; x, y, z) = \sum_{i,j=0}^{\infty} \frac{(-1)^{i+j} (\alpha)_{i+j} (\beta)_{i+j} (\gamma_1 - \varepsilon_1)_i (\gamma_2 - \varepsilon_2)_j}{(\gamma_1)_i (\gamma_2)_j (\varepsilon_1)_i (\varepsilon_2)_j i! j!} x^i y^j \times F_C(\alpha + i + j, \beta + i + j; \varepsilon_1 + i, \varepsilon_2 + j, \gamma_3; x, y, z), \quad (3.48)$$

$$\begin{aligned}
& F_C(\alpha, \beta; \gamma_1, \gamma_2, \gamma_3; x, y, z) \\
&= \sum_{i,j,k=0}^{\infty} \frac{(-1)^{i+j+k} (\alpha)_{i+j+k} (\beta)_{i+j+k} (\gamma_1 - \varepsilon_1)_i (\gamma_2 - \varepsilon_2)_j (\gamma_3 - \varepsilon_3)_k}{(\gamma_1)_i (\gamma_2)_j (\gamma_3)_k (\varepsilon_1)_i (\varepsilon_2)_j (\varepsilon_3)_k i! j! k!} x^i y^j z^k \\
&\quad \times F_C(\alpha + i + j + k, \beta + i + j + k; \varepsilon_1 + i, \varepsilon_2 + j, \varepsilon_3 + k; x, y, z),
\end{aligned} \tag{3.49}$$

$$F_D(\alpha; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = \sum_{i=0}^{\infty} \frac{(-1)^i (\alpha)_i (\varepsilon_1 - \beta_1)_i}{(\gamma)_i i!} x^i F_D(\alpha + i; \varepsilon_1 + i, \beta_2, \beta_3; \gamma + i; x, y, z), \tag{3.50}$$

$$F_D(\alpha; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = \sum_{i=0}^{\infty} \frac{(\alpha)_i (\beta_1 - \varepsilon_1)_i}{(\gamma)_i i!} x^i F_D(\alpha + i; \varepsilon_1, \beta_2, \beta_3; \gamma + i; x, y, z), \tag{3.51}$$

$$\begin{aligned}
& F_D(\alpha; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) \\
&= \sum_{i,j=0}^{\infty} \frac{(-1)^{i+j} (\alpha)_{i+j} (\varepsilon_1 - \beta_1)_i (\varepsilon_2 - \beta_2)_j}{(\gamma)_{i+j} i! j!} x^i y^j F_D(\alpha + i + j; \varepsilon_1 + i, \varepsilon_2 + j, \beta_3; \gamma + i + j; x, y, z),
\end{aligned} \tag{3.52}$$

$$\begin{aligned}
& F_D(\alpha; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) \\
&= \sum_{i,j=0}^{\infty} \frac{(\alpha)_{i+j} (\beta_1 - \varepsilon_1)_i (\beta_2 - \varepsilon_2)_j}{(\gamma)_{i+j} i! j!} x^i y^j F_D(\alpha + i + j; \varepsilon_1, \varepsilon_2, \beta_3; \gamma + i + j; x, y, z),
\end{aligned} \tag{3.53}$$

$$\begin{aligned}
& F_D(\alpha; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = \sum_{i,j,k=0}^{\infty} \frac{(-1)^{i+j+k} (\alpha)_{i+j+k} (\varepsilon_1 - \beta_1)_i (\varepsilon_2 - \beta_2)_j (\varepsilon_3 - \beta_3)_k}{(\gamma)_{i+j+k} i! j! k!} x^i y^j z^k \\
&\quad \times F_D(\alpha + i + j + k; \varepsilon_1 + i, \varepsilon_2 + j, \varepsilon_3 + k; \gamma + i + j + k; x, y, z),
\end{aligned} \tag{3.54}$$

$$\begin{aligned}
& F_D(\alpha; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) = \sum_{i,j,k=0}^{\infty} \frac{(\alpha)_{i+j+k} (\beta_1 - \varepsilon_1)_i (\beta_2 - \varepsilon_2)_j (\beta_3 - \varepsilon_3)_k}{(\gamma)_{i+j+k} i! j! k!} x^i y^j z^k \\
&\quad \times F_D(\alpha + i + j + k; \varepsilon_1, \varepsilon_2, \varepsilon_3; \gamma + i + j + k; x, y, z).
\end{aligned} \tag{3.55}$$

4 Demonstrations of some the decompositions formulas for Lauricella functions in the three-variable cases

The various decomposition formulas for the Lauricella functions F_A , F_B , F_C and F_D in three variables (which are presented here and in other places in the previously cited literature) can be proven fairly simply by suitably applying superposition of the inverse pairs of symbolic operators introduced in Section 1. As an example, we shall briefly indicate the proof of the decomposition formula (3.54). For the three-variable Lauricella function F_D , it is not difficult to show from (3.26) that

$$\begin{aligned}
& F_D(\alpha; \beta_1, \beta_2, \beta_3; \gamma; x, y, z) \\
&= \sum_{i=0}^{\infty} \frac{(\varepsilon_1 - \beta_1)_i (-\delta_1)_i}{(\varepsilon_1)_i i!} \sum_{j=0}^{\infty} \frac{(\varepsilon_2 - \beta_2)_j (-\delta_2)_j}{(\varepsilon_2)_j j!} \sum_{k=0}^{\infty} \frac{(\varepsilon_3 - \beta_3)_k (-\delta_3)_k}{(\varepsilon_3)_k k!} F_D(\alpha; \varepsilon_1, \varepsilon_2, \varepsilon_3; \gamma; x, y, z),
\end{aligned} \tag{4.1}$$

$$\left(\delta_1 := x \frac{\partial}{\partial x}; \delta_2 := y \frac{\partial}{\partial y}; \delta_3 := z \frac{\partial}{\partial z} \right).$$

Furthermore, by a straightforward computation, we have

$$(-\delta_1)_i F_D(\alpha; \varepsilon_1, \varepsilon_2, \varepsilon_3; \gamma; x, y, z) = (-1)^i x^i \frac{(\alpha)_i (\varepsilon_1)_i}{(\gamma)_i} F_D(\alpha + i; \varepsilon_1 + i, \varepsilon_2, \varepsilon_3; \gamma + i; x, y, z), \tag{4.2}$$

$$\begin{aligned}
& (-\delta_2)_j (-\delta_1)_i F_D(\alpha; \varepsilon_1, \varepsilon_2, \varepsilon_3; \gamma; x, y, z) \\
&= (-1)^{i+j} x^i y^j \frac{(\alpha)_{i+j} (\varepsilon_1)_i (\varepsilon_2)_j}{(\gamma)_{i+j}} F_D(\alpha + i + j; \varepsilon_1 + i, \varepsilon_2 + j, \varepsilon_3; \gamma + i + j; x, y, z)
\end{aligned} \tag{4.3}$$

and

$$\begin{aligned}
& (-\delta_3)_k (-\delta_2)_j (-\delta_1)_i F_D(\alpha; \varepsilon_1, \varepsilon_2, \varepsilon_3; \gamma; x, y, z) = (-1)^{i+j+k} x^i y^j z^k \\
& \times \frac{(\alpha)_{i+j+k} (\varepsilon_1)_i (\varepsilon_2)_j (\varepsilon_3)_k}{(\gamma)_{i+j+k}} F_D(\alpha + i + j + k; \varepsilon_1 + i, \varepsilon_2 + j, \varepsilon_3 + k; \gamma + i + j + k; x, y, z).
\end{aligned} \tag{4.4}$$

Upon substituting from (4.4) into (4.1), we finally arrive at the decomposition formula (3.54).

5 Integral representations via decomposition formulas

Here in this section, we observe that several integral representations of the Eulerian type can be deduced also from the corresponding decomposition formulas of Section 3. For example, using integral representation ([1], p. 115, (5))

$$\begin{aligned}
F_A^{(r)}(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) &= \frac{\Gamma(\gamma_1) \Gamma(\gamma_2) \Gamma(\gamma_3)}{\Gamma(\beta_1) \Gamma(\beta_2) \Gamma(\beta_3) \Gamma(\gamma_1 - \beta_1) \Gamma(\gamma_2 - \beta_2) \Gamma(\gamma_3 - \beta_3)} \\
& \int_0^1 \dots \int_0^1 t_1^{\beta_1-1} t_2^{\beta_2-1} t_3^{\beta_3-1} (1-t_1)^{\gamma_1-\beta_1-1} (1-t_2)^{\gamma_2-\beta_2-1} (1-t_3)^{\gamma_3-\beta_3-1} (1-xt_1-yt_1-zt_3)^{-\alpha} dt_1 dt_2 dt_3, \\
& \text{Re } \gamma_l > \text{Re } \beta_l > 0, \quad l = 1, 2, 3
\end{aligned} \tag{5.1}$$

we define

$$\begin{aligned}
F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) &= \frac{\Gamma(\gamma_1) \Gamma(\gamma_2) \Gamma(\gamma_3)}{\Gamma(\varepsilon_1) \Gamma(\varepsilon_2) \Gamma(\varepsilon_3) \Gamma(\gamma_1 - \varepsilon_1) \Gamma(\gamma_2 - \varepsilon_2) \Gamma(\gamma_3 - \varepsilon_3)} \\
& \int_0^1 \dots \int_0^1 t_1^{\varepsilon_1-1} t_2^{\varepsilon_2-1} t_3^{\varepsilon_3-1} (1-t_1)^{\gamma_1-\varepsilon_1-1} (1-t_2)^{\gamma_2-\varepsilon_2-1} (1-t_3)^{\gamma_3-\varepsilon_3-1} \\
& \times (1-yt_2-zt_3)^{-\alpha} F\left(\alpha, \beta_1; \varepsilon_1; \frac{xt_1}{1-yt_2-zt_3}\right) dt_1 dt_2 dt_3, \\
& \text{Re } \gamma_1 > \text{Re } \varepsilon_1 > 0, \quad \text{Re } \gamma_l > \text{Re } \beta_l > 0, \quad l = 2, 3
\end{aligned} \tag{5.2}$$

$$\begin{aligned}
F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) &= \frac{\Gamma(\gamma_1) \Gamma(\gamma_2) \Gamma(\gamma_3)}{\Gamma(\varepsilon_1) \Gamma(\varepsilon_2) \Gamma(\varepsilon_3) \Gamma(\gamma_1 - \varepsilon_1) \Gamma(\gamma_2 - \varepsilon_2) \Gamma(\gamma_3 - \varepsilon_3)} \\
& \times \int_0^1 \dots \int_0^1 t_1^{\varepsilon_1-1} t_2^{\varepsilon_2-1} t_3^{\varepsilon_3-1} (1-t_1)^{\gamma_1-\varepsilon_1-1} (1-t_2)^{\gamma_2-\varepsilon_2-1} (1-t_3)^{\gamma_3-\varepsilon_3-1} (1-xt_1-yt_1-zt_3)^{-\alpha} \\
& \times F_2\left(\alpha; \beta_1 - \varepsilon_1, \beta_2 - \varepsilon_2; \gamma_1 - \varepsilon_1, \gamma_2 - \varepsilon_2; \frac{x(1-t_1)}{1-xt_1-yt_1-zt_3}, \frac{y(1-t_2)}{1-xt_1-yt_1-zt_3}\right) dt_1 dt_2 dt_3 \\
& \text{Re } \gamma_l > \text{Re } \varepsilon_l > 0, \quad l = 1, 2, \quad \text{Re } \gamma_3 > \text{Re } \beta_3 > 0,
\end{aligned} \tag{5.3}$$

$$\begin{aligned}
F_A(\alpha; \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_3; x, y, z) &= \frac{\Gamma(\gamma_1) \Gamma(\gamma_2) \Gamma(\gamma_3)}{\Gamma(\varepsilon_1) \Gamma(\varepsilon_2) \Gamma(\varepsilon_3) \Gamma(\gamma_1 - \varepsilon_1) \Gamma(\gamma_2 - \varepsilon_2) \Gamma(\gamma_3 - \varepsilon_3)} \\
&\times \int_0^1 \dots \int_0^1 t_1^{\varepsilon_1-1} t_2^{\varepsilon_2-1} t_3^{\varepsilon_3-1} (1-t_1)^{\gamma_1-\varepsilon_1-1} (1-t_2)^{\gamma_2-\varepsilon_2-1} (1-t_3)^{\gamma_3-\varepsilon_3-1} (1-xt_1-yt_1-zt_3)^{-\alpha} \\
&\times F_A(\alpha; \beta_1 - \varepsilon_1, \beta_2 - \varepsilon_2, \beta_3 - \varepsilon_3; \gamma_1 - \varepsilon_1, \gamma_2 - \varepsilon_2, \gamma_3 - \varepsilon_3; \\
&\quad \frac{x(1-t_1)}{1-xt_1-yt_1-zt_3}, \frac{y(1-t_2)}{1-xt_1-yt_1-zt_3}, \frac{z(1-t_3)}{1-xt_1-yt_1-zt_3}) dt_1 dt_2 dt_3, \\
&\text{Re } \gamma_l > \text{Re } \varepsilon_l > 0, \quad l = 1, 2, 3.
\end{aligned} \tag{5.4}$$

Remark. Introduced operators can be used for other hypergeometric functions also.

References

- [1] P. Appell and J. Kampe de Fériet, *Fonctions Hypergeometriques et Hyperspheriques; Polynomes d'Hermite*, Gauthier - Villars. Paris, 1926.
- [2] L. Bers, *Mathematical Aspects of Subsonic and Transonic Gas Dynamics*, Wiley, New York, 1958.
- [3] J.L. Burchnall, T.W. Chaundy, Expansions of Appell's double hypergeometric functions, *Quart. J. Math.* Oxford Ser. 11 (1940), 249-270.
- [4] J.L. Burchnall, T.W. Chaundy, Expansions of Appell's double hypergeometric functions. II, *Quart. J. Math.* Oxford Ser. 12 (1941), 112-128.
- [5] T.W. Chaundy, Expansions of hypergeometric functions, *Quart. J. Math.* Oxford Ser. 13, (1942), 159-171.
- [6] F.I. Frankl, *Selected Works in Gas Dynamics*. Nauka, Moscow 1973 (in Russian).
- [7] A. Hasanov, Fundamental solutions of generalized bi-axially symmetric Helmholtz equation. *Complex Variables and Elliptic Equations*. 52 (8) (2007) 673-683.
- [8] A. Hasanov, The solution of the Cauchy problem for generalized Euler-Poisson-Darboux equation. *International Journal of Applied Mathematics and Statistics*. 8 (7) (2007) 30-44.
- [9] A. Hasanov, On a mixed problem for the equation $\text{sign } y |y|^m u_{xx} + x^n u_{yy} = 0$. *Izv. Akad. Nauk UzSSR*, ser. Fiz.-mat. Nauk, 2 (1982) 28-32. (in Russian)
- [10] A. Hasanov and H. M. Srivastava, Some decomposition formulas associated with the Lauricella function $F_A^{(r)}$ and other multiple hypergeometric functions, *App. Math. Lett.* 19 (2006) 113-121.
- [11] A. Hasanov and H.M. Srivastava, Decomposition Formulas Associated with the Lauricella Multi-variable Hypergeometric Functions. *Computers and Mathematics with Applications*, 53 (7) (2007) 1119-1128.

- [12] G. Lohofer, Theory of an electromagnetically deviated metal sphere. 1: Absorbed power. *SIAM J. Appl. Math.* 49 (1989) 567-581.
- [13] A.M. Mathai, R. K. Saxena, *Generalized Hypergeometric Functions with Applications in Statistics and Physical Sciences*. Springer-Verlag, Berlin, Heidelberg and New York. 1973.
- [14] O.I. Marichev, *Handbook of Integral Transforms of Higher Transcendental Functions: Theory and algorithmic Tables*. Halsted Press (Ellis Horwood Limited, Chichester), Wiley, New York, Chichester, Brisbane and Toronto, 1982.
- [15] A.W. Niukkanen, Generalised hypergeometric series ${}^N F(x_1, \dots, x_N)$ arising in physical and quantum chemical applications, *J. Phys. A: Math. Gen.* 16 (1983) 1813-1825.
- [16] S.B. Opps, N. Saad, H.M. Srivastava, Some reduction and transformation formulas for the Appell hypergeometric function F_2 , *J. Math. Anal. Appl.* 302 (2005) 180-195.
- [17] P.A. Padmanabham, H.M. Srivastava, Summation formulas associated with the Lauricella function $F_A^{(r)}$, *Appl. Math. Lett.* 13 (1) (2000) 65-70.
- [18] E.G. Poole, *Introduction to the Theory of Linear Differential Equations*, Clarendon (Oxford University Press), Oxford, 1936.
- [19] I.N. Sneddon. *Special Functions of Mathematical Physics and Chemistry*. Third ed., Longman, London and New York. 1980.
- [20] H.M. Srivastava., B.R.K. Kashyap. *Special Functions in Queuing Theory and Related Stochastic Processes*. Academic Prees, New York, London and San Francisco, 1982.
- [21] H.M. Srivastava, P.W. Karlsson, *Multiple Gaussian Hypergeometric Series*, Halsted Press (Ellis Horwood Limited, Chichester), Wiley, New York, Chichester, Brisbane and Toronto, 1985.